Sanum

Latensin® 6X Suppositories

Registration:	No registration in Germany
Administration form:	Suppositories for rectal application
Preparation group:	Bacterial preparation
Active ingredient:	Bacillus cereus e volumine ex muris cellulae (lyophil., steril.) 6X
Composition:	1 suppository contains: Medically active substance: 0.2 g Bacillus cereus e volumine ex muris cellulae (lyophil., steril.) 6X trit. Other constituents: hard fat, lactose.
According to experience, to be administered with caution:	As with all registered homeopathic remedies, therapeutic indications are not stated.
Application:	Unless otherwise prescribed: 1-3 x weekly insert 1 suppository rectally.
Side effects:	Because of specific organic components of Latensin®, theoretically, hypersensitivity may occur. In this case, discontinue medication and treat symptomatically.
Contraindications:	Fever, conditions of weakness. Do not administer in cases of known hypersensitivity to Bacillus cereus.
Adverse reactions:	None known. Generally well tolerated. In the first 2 or 3 days after the application sometimes conditions of weakness, fortified expectoration or subfebrile temperatures may show. Strong general reactions are always a sign of overdose.

Interactions with other remedies:	Immunosuppressive drugs can influence the effectivity of Latensin®. An interval of 4 weeks before and after treatment with orally administered live vaccines must be observed.
Precautions:	As with all medications and due to the variations of clinical studies, professional medical advice should be sought prior to recommending this product to women during pregnancy or breastfeeding, as well as with children.
Advice:	This product contains lactose.
Duration of treatment:	Dependent on the advice of the physician or health care professional.
How supplied:	The following dosage forms are available: 5 ml dropper bottle 6X, 1 ml ampule 10 and 50 4X, 6X, 10 suppositories 6X, 5 capsules 4X, 6X.
This product is availabe in:	- Italien - Schweiz - Niederlande